Simplicial Complexes

Lecture 3 - CMSE 890

Prof. Elizabeth Munch

Michigan State University

Dept of Computational Mathematics, Science & Engineering


Tues, Sep 2, 2025

Goals

Goals for today: Ch 2.1

- Define geometric/abstract simplicial complexes
- Define simplicial maps

A simplicial complex is a generalizion of a graph



General position

A set $\{a_0, \dots, a_n\} \subset \mathbb{R}^N$ is **in general position** if no k of them lie in a k-2 dimensional hyperplane (k = 2, 3, ..., N + 1).

Not on the same line (1-din hyperplane)

they are in gen pos.

Warning: Papers will often modify this definition to suit their purposes!

n-Plane

Given a set of points $\{a_0, \dots, a_n\}$ in general position, the *n*-plane *P* spanned by the points is

$$P = \left\{ \sum_{i=0}^{n} t_{i} a_{i} \in \mathbb{R}^{N} \mid \sum t_{i} = 1 \right\}.$$

$$\left\{ a_{0}, a_{1} \right\}$$

$$t_{0} = 1$$

Liz Munch (MSU-CMSE)

Lec 3


Tues, Sep 2, 2025


n-simplex

Given a set of points $\{a_0, \dots, a_n\}$ in general position, the *n*-simplex σ spanned by the points is

$$P = \left\{ \sum_{i=0}^n t_i a_i \in \mathbb{R}^N \mid \sum t_i = 1
ight\}$$


such that $t_i \geq 0 \ \forall i$.





Barycentric coordinates

The numbers t_i are uniquely determined t_i and are called **barycentric coordinates**.




More definitions

- $\{a_0, \dots, a_n\}$ are called the **vertices** of σ .
- The dimension of $\sigma = [a_0, \dots, a_n]$ is n.
- Any simplex spanned by a subset of $\{a_0, \dots, a_n\}$ is a **face** of σ .
- A face is proper if it isn't the simplex itself.


TKT


"18 a face of "



Even more definitions

- The union of the proper faces is the **boundary** of σ , $Bd(\sigma)$.
- The **interior** of σ is $\sigma Bd(\sigma)$. This is sometimes called the open simplex.







Geometric simplicial complex


A (geometric) simplicial complex K in \mathbb{R}^N is a (finite) collection of simplices in \mathbb{R}^N such that

• Every face of a simplex of K is in K.

 $oldsymbol{\circ}$ The intersection of any two simplices of K is a face of each.










Dimension


The **dimension** of K is the maximum dimension of its simplices.




$$d_{im}(K) = 2$$

Abstract simplicial complexes

An abstract simplicial complex K is a (finite) collection of (finite) non-empty sets of V such that $\underline{\alpha} \in K$ and $\beta \subseteq \alpha$ implies $\beta \in K$.



Two types




Liz Munch (MSU-CMSE)

Lec 3

Tues, Sep 2, 2025

Subcomplex


If L is a subcollection of K that contains all faces of its elements, then L is called a **subcomplex**.



A subcomplex $L \subset K$ is **full** if it contains all simplices of K spanned by the vertices in L.

Skeleton

The subcomplex of K consisting of all simplices of dimension $\leq p$ is called the p-skeleton of K, usually denoted K^p .



$$St(\sigma) = \{ \tau \in K \mid \sigma \in \tau \}$$




• The star of a simplex $\sigma \in K$, $\operatorname{St}(\sigma)$, is the union of the interiors of simplices in K which

have σ as a face. Warning: $\operatorname{St}(\sigma) \subset |K|$ is not a simplicial complex!


- The closed star, $\overline{\mathrm{St}}(\sigma)$, is the closure of $\mathrm{St}(\sigma)$.
- · Line St(r) St(r)

 $T = \{c_1b\}$ $T_1 = \{a_1b_1c_3\}$ $T_2 = \{b_1c_1d\}$




Links

The **link** of a simplex is $\overline{\mathrm{St}}(\sigma) - \mathrm{St}(\sigma)$.



Given a finite collection of sets \mathcal{F} , the **nerve** is


$$Nrv(\mathcal{U}) = \{X \subseteq \mathcal{F} \mid \bigcap_{U \in X} U \neq \emptyset\}.$$



Check: This is actually an abstract simplicial complex!

Recall: K is a simplicial complex if $\sigma \in K$ and $\tau \leq \sigma$ implies $\tau \in K$.


Try it: What is the nerve of this collection of disks?



Try it: What is the nerve of this collection of sets?



The difference between the examples





Homework

Pick one

- Let *K* be a simplicial complex. Show that *K* is the nerve of the collection of stars of its vertices.
- A flag in a simplicial complex K is a nested sequence of proper faces σ₀ < σ₁ < ··· < σ_k.
 Show that the collection of flags in K forms a simplicial complex; this is called the **order complex** of K.

For either question, show the construction on the simplicial complex below.

